

**CHARACTERIZATION OF OZONE EPISODES IN THE SOUTH COAST  
AIR BASIN: EFFECTS OF AIR PARCEL RESIDENCE TIME AND  
WEEKEND/WEEKDAY DIFFERENCES**

Final Report  
Contract No. 93-316  
California Air Resources Board

**Principal Investigators**

Warren Blier

Department of Atmospheric Sciences  
College of Letters and Sciences  
University of California, Los Angeles

and

Arthur M. Winer

Environmental Science and Engineering Program  
Department of Environmental Health Sciences  
School of Public Health  
University of California, Los Angeles

**Graduate Student Researchers**

Darin Hansen  
Namita Verma  
University of California, Los Angeles

**June 28, 1996**

## ABSTRACT

To date, airshed modeling for control strategy evaluations has typically involved simulations of only a few air pollution episodes, which may not represent either the complete range of meteorological conditions over which high ozone episodes can occur, or differences in ozone concentrations resulting from variation in precursor emissions by day-of-the-week. The present project focused on the California South Coast Air Basin (SoCAB) "smog season" of 1 May to 31 October for the years 1986-1993, and had the following primary objectives: to better characterize the range of meteorological conditions associated with high ozone episodes in the SoCAB; to assess the variation in, and influence of, transport and residence time on the occurrence of high concentrations of ozone; to isolate and examine relationships between day-of-the-week and ozone levels; and to examine sub-regional air quality trends over the past decade, including degree of correlation between ozone and NO<sub>2</sub> ambient air concentrations. Fulfillment of these research objectives necessitated the development of comprehensive, critically evaluated air quality and meteorological databases for the SoCAB. In addition to their utility for the present research project, these databases should be of benefit to future air quality research concerning the SoCAB.

Results of this project suggest the need for further investigation of the longstanding view of the importance of air parcel transport across the SoCAB. Specifically, better correlations between NO<sub>2</sub> and ozone concentrations were observed within sub-regions than between them. This finding (while not establishing causality) is consistent with the fact that although the Coastal and Metropolitan subregions were formerly much more densely populated than other regions of the Basin, over the past two decades the major growth in the SoCAB has occurred in the middle and eastern parts of the Basin, producing a commensurate change in emission patterns.

These results are also consistent with those from a comparative analysis of the surface meteorological conditions associated with the days of highest SoCAB peak ozone and days associated with average peak-ozone concentrations. Examination of the vector-mean hourly-average resultant winds throughout the SoCAB for each hour of the day for the high ozone days, along with an analysis of the average time at which the peak ozone value was reported at each station, indicated surface transport on the time scale of a day was of a subregional scale. (However, determination of the actual transport would require wind data aloft which were not available in the present project.) Also of interest was that both average- and highest-ozone days were associated with very similar vector-mean hourly-average resultant winds throughout the SoCAB. Thus, it did not appear differences in transport, at least as far as this was indicated by the surface winds, played a generally significant role in determining the magnitude of the Basin-peak ozone value.

Significant relationships were found, however, between Basin-peak ozone values and both the 850 mb temperature, and the Basin-wide values of maximum surface air temperature. Consistent with earlier studies, temperatures tended to be significantly higher on the high ozone days than on the middle ozone days. Some relationship was also found between the characteristic form of the synoptic-scale weather pattern and corresponding ozone concentrations: specifically, high ozone days tended to occur more often in association with

an "established ridge" aloft (characterized by high mid-tropospheric geopotential heights, a warm lower- and mid-tropospheric air mass, and light winds aloft), while a synoptic-scale pattern with lower heights and stronger and more zonal flow aloft was more typical for middle-ozone days.

Trend analyses were performed by comparing the mean Basin-maximum ozone values of the various CART nodes for 1986-1989 with those for 1990-1993; results indicated that regardless of the meteorological conditions, generally lower peak ozone values were observed in the latter four years than in the first four years.

A further trend analysis was performed by examining the means of the ten-highest hourly-average concentrations for each of the four-year periods. These "worst ozone days" trends showed the most pronounced percentage decrease in these highest ozone concentrations occurred in the western or middle portion of the Basin, corresponding generally to the area of maximum percentage decrease in early morning  $\text{NO}_x$  ambient concentrations for these same highest ozone days.

Both cumulative hourly ozone exposure above the federal standard of 12 pphm and number of first stage alerts were also found to decrease over the eight-year period of interest for each day of the week for the stations at Central Los Angeles, Azusa, and Riverside (chosen as representative of the western, middle and eastern portions of the urbanized area of the Basin). Higher daily peak ozone concentrations remained more common on weekend days, though, while lower daily maximum concentrations occurred more frequently on weekdays.

Although there was a lack of evidence for  $\text{NO}_2$  or  $\text{NO}_x$  carry-over influencing next-day peak ozone, further investigation of the effect of carry-over of weekday precursor emissions on weekend high ozone episodes is needed. Better correlation of peak ozone with morning  $\text{NO}_2$  than with morning  $\text{NO}_x$  ambient air concentrations was observed.

Implications of all of these results for  $\text{NO}_x$  and VOC control strategies will remain unclear until more accurate emissions data become available for the SoCAB as a function of day of the week, and by sub-region.

## ACKNOWLEDGEMENTS

The contributions of several members of the California Air Resources Board (ARB) staff, particularly Ash Lashgari and Bart Croes of the Research Division and Larry Larsen of the Technical Support Division, were greatly appreciated.

We wish to thank Dennis King and Leon Dolislager of the ARB for providing air quality data and assisting in interpreting the station and variable codes. Dennis King also provided copies of several ARB reports and documents. We express our appreciation to Joseph Cassmassi of the SCAQMD for providing meteorological and air quality data and for valuable discussions regarding the CART analysis in this study. Eric Fujita at the Desert Research Institute provided us with assistance in interpreting early morning NO<sub>2</sub> and NO<sub>x</sub> data. Thanks also to Dale Shimp of the ARB for providing us with information on emission inventories.

We wish to thank Martin Wachs, UCLA Department of Urban Planning, who provided us with information about transportation studies, and Nick Jones from the Los Angeles office of Caltrans for providing us with traffic count data. We are grateful to Qunfei Ma, UCLA Department of Atmospheric Sciences, for helping complete the upper-air database, and especially thank James Murakami, of the same department, for providing access to many archived air quality and meteorological data sets. Thanks also to Kevin Durkee at the SCAQMD who assisted us by providing resultant wind data and Fabrice Cuq for substantial assistance with data file management and computer software. We are grateful to James Adams and Teri McLaughlin for their extensive and thorough assistance in the assembly and production of this report.

This report was submitted in fulfillment of Contract No. 93-316. Work on this project was completed on February 27, 1996.

## **DISCLAIMER**

The statements and conclusions in this report are those of the contractors and not necessarily those of the California Air Resources Board. The mention of commercial products, their source, or their use in connection with the material reported herein, is not to be construed as either an actual or implied endorsement of such products.

# TABLE OF CONTENTS

	<u>Page</u>
Abstract .....	i
Acknowledgements .....	iii
Disclaimer .....	iv
List of Tables .....	ix
List of Figures .....	xiii
Glossary of Terms, Abbreviations, and Symbols .....	xix
 1.0 EXECUTIVE SUMMARY .....	 1-1
1.1 Introduction .....	1-1
1.2 Objectives .....	1-2
1.3 Database Development .....	1-3
1.4 Correlations of Ozone Concentrations With NO <sub>2</sub> and NO <sub>x</sub> Ambient Concentrations as a Function of Day of Week and Subregion .....	 1-4
1.4.1 Daily Basin O <sub>3</sub> Maximum Concentrations vs. Morning NO <sub>2</sub> and NO <sub>x</sub> Concentrations .....	 1-6
1.4.2 Daily Subregion O <sub>3</sub> Maximum Concentrations vs. Morning NO <sub>2</sub> and NO <sub>x</sub> Concentrations .....	 1-6
1.4.3 Daily O <sub>3</sub> Maximum vs. Previous Evening's NO <sub>2</sub> and NO <sub>x</sub> Concentrations .....	 1-8
1.5 Examination of Air Parcel Residence Time and Other Meteorological Parameters .....	 1-10
1.5.1 Surface Meteorological Conditions: High Ozone Days vs. Middle Ozone Days .....	 1-11
1.5.2 850 mb Meteorological Conditions: High Ozone Days vs. Middle Ozone Days .....	 1-13
1.5.3 Pattern Number: High Ozone Days vs. Middle Ozone Days ...	1-14
1.5.4 Application of the CART Scheme .....	1-15
1.6 Characterization of Weekday/Weekend Effects .....	1-16
1.6.1 Daily Maximum Ozone Distributions for Weekdays vs. Weekend Days .....	 1-16
1.6.2 Ozone First Stage Alerts and Cumulative Hourly Exceedance for Weekdays vs. Weekend Days .....	 1-16
1.6.3 Weekday/Weekend Analyses Using the CART Scheme .....	1-17
1.6.4 Examination of Ten Days with Highest Hourly-Average Ozone Concentrations: 1986-1989 vs. 1990-1993 .....	 1-18
1.6.5 Weekday/Weekend Differences in Ambient NO <sub>2</sub> and NO <sub>x</sub> Concentrations and NO <sub>2</sub> /NO <sub>x</sub> Ratios .....	 1-18

1.7 Principal Findings	1-21
2.0 INTRODUCTION, BACKGROUND AND OBJECTIVES	2-1
2.1 Introduction	2-1
2.2 Background	2-3
2.3 Objectives	2-5
2.3.1 Overall Objectives	2-5
2.3.2 Specific Objectives	2-5
3.0 ACQUISITION, DEVELOPMENT AND VALIDATION OF DATABASES	3-1
3.1 Database Acquisition	3-1
3.1.1 Air Quality Database	3-1
3.1.2 Meteorological Database	3-2
3.2 Comparison of ARB and SCAQMD Databases	3-2
3.3 Database Completeness	3-2
3.3.1 Initial Completeness Analysis	3-2
3.3.2 Additional Completeness Tests	3-8
4.0 CORRELATIONS OF OZONE CONCENTRATIONS WITH NO <sub>2</sub> AND NO <sub>x</sub> AMBIENT CONCENTRATIONS AS A FUNCTION OF DAY OF WEEK AND SUBREGION	4-1
4.1 Daily Basin O <sub>3</sub> Maximum Concentrations vs. Morning NO <sub>2</sub> and NO <sub>x</sub> Concentrations	4-4
4.1.1 Coastal and Metropolitan Subregions	4-4
4.1.2 San Gabriel Valley Subregion	4-5
4.1.3 Inland Valley Subregion	4-6
4.2 Daily Subregion O <sub>3</sub> Maximum Concentrations vs. Morning NO <sub>2</sub> and NO <sub>x</sub> Concentrations	4-9
4.2.1 Coastal and Metropolitan Subregions	4-9
4.2.2 San Gabriel Valley Subregion	4-10
4.2.3 Inland Valley Subregion	4-11
4.3 Daily O <sub>3</sub> Maximum vs. Previous Evening's NO <sub>2</sub> and NO <sub>x</sub> Concentrations	4-13
4.3.1 Coastal and Metropolitan Subregions	4-14
4.3.2 San Gabriel Valley Subregion	4-15
4.4 Summary	4-16

5.0 EXAMINATION OF AIR PARCEL RESIDENCE TIME AND OTHER METEOROLOGICAL PARAMETERS .....	5-1
5.1 Surface Meteorological Conditions: High Ozone Days vs. Middle Ozone Days .....	5-1
5.1.1 Resultant Surface Winds .....	5-2
5.1.2 Time of Peak Ozone Occurrence .....	5-7
5.1.3 Backward Surface Trajectories .....	5-19
5.1.4 Surface Wind Roses .....	5-21
5.1.5 Surface Temperatures .....	5-27
5.2 850 mb Meteorological Conditions: High Ozone Days vs. Middle Ozone Days .....	5-27
5.2.1 850 mb Temperatures .....	5-31
5.2.2 850 mb Winds .....	5-32
5.2.3 850 mb Wind Roses .....	5-35
5.3 Pattern Number: High Ozone Days vs. Middle Ozone Days .....	5-37
5.4 Application of the CART Scheme .....	5-39
5.4.1 Individual Years: 1986-93 .....	5-43
5.4.2 1986-89 vs. 1990-93 .....	5-48
5.4.3 High Ozone Days vs. Middle Ozone Days .....	5-48
5.5 Summary .....	5-50
6.0 OZONE TRENDS AND WEEKDAY/WEEKEND EFFECTS .....	6-1
6.1 Introduction .....	6-1
6.2 Seasonal Mean Daily Maximum Ozone By Day of the Week .....	6-1
6.3 Daily Maximum Ozone Distributions for Weekdays vs. Weekends Days .....	6-2
6.4 Ozone First Stage Alerts and Cumulative Hourly Exposure for Weekdays vs. Weekends Days .....	6-28
6.5 Weekday/Weekend Analyses Using the CART Scheme .....	6-40
6.6 Examination of Ten Days with Highest Hourly-Average Ozone Concentrations: 1986-89 vs. 1990-93 .....	6-42
6.6.1 All Days .....	6-42
6.6.2 Weekdays and Weekend Days: Ozone Trends Between 1986-89 and 1990-93 .....	6-44
6.6.3 Weekdays vs. Weekend Days: Ozone Differences Within Periods .....	6-49
6.7 Examination of Individual Days with Highest Hourly-Average	



Ozone Concentrations for Each Year in the Period 1986-93	.... 6-49
6.8 NO <sub>2</sub> /NO <sub>x</sub> Trends and Weekday/Weekend Differences in Ambient NO <sub>2</sub> and NO <sub>x</sub> Concentrations, and NO <sub>2</sub> /NO <sub>x</sub> Ratios	.... 6-54
6.8.1 Previous Studies	..... 6-54
6.8.2 Calculation of Mean Morning NO <sub>2</sub> and NO <sub>x</sub> Concentrations and NO <sub>2</sub> /NO <sub>x</sub> Ratios	..... 6-56
6.8.3 Statistical Significance of Observed Differences	..... 6-59
6.8.4 NO <sub>2</sub> and NO <sub>x</sub> Trends for Ten Highest Ozone Days	..... 6-59
6.8.5 NO <sub>2</sub> Concentrations for 28 Highest and Middle Ozone Days	.. 6-59
6.9 Summary	..... 6-65
7.0 EXPLORATORY STUDIES	..... 7-1
7.1 Meteorological Representativeness of the August 1987 SCAQS Episode	. 7-2
7.1.1 Resultant Surface Winds	..... 7-2
7.1.2 Time of Peak Ozone Occurrence	..... 7-4
7.1.3 Surface Temperatures	..... 7-8
7.1.4 850 mb Temperatures	..... 7-8
7.1.5 850 mb Winds	..... 7-8
7.1.6 Pattern Number	..... 7-10
7.1.7 CART Scheme Classification	..... 7-10
7.2 Multiple Linear Regression Relationships Between Meteorological Parameters and Ozone	..... 7-11
8.0 SUMMARY AND CONCLUSIONS	..... 8-1
8.1 Correlation of Ozone Concentrations with NO <sub>2</sub> and NO <sub>x</sub> Ambient Concentrations	..... 8-1
8.2 Meteorology Analyses	..... 8-1
8.3 Ozone Trends and Weekday/Weekend Effects	..... 8-2
8.4 Limitations to the Present Analysis	..... 8-3
9.0 RECOMMENDATIONS FOR FUTURE RESEARCH	..... 9-1
10.0 REFERENCES	..... 10-1



## LIST OF TABLES

<u>Table Number</u>	<u>Title</u>	<u>Page</u>
1-1	Air quality and meteorological databases (1986-93) acquired for the South Coast Air Basin .....	1-3
1-2	Subregions of the SoCAB employed in this study and air monitoring stations within each subregion for which air quality data were employed .....	1-5
1-3	Correlation coefficients between daily Coastal/Metropolitan ozone maximum concentration and average morning NO <sub>2</sub> and NO <sub>x</sub> concentrations in the Coastal/Metropolitan subregion .....	1-7
1-4	Correlation coefficients between daily Inland Valley ozone maximum concentration and average morning NO <sub>2</sub> and NO <sub>x</sub> concentrations in the Inland Valley subregion .....	1-8
3-1	The air quality and meteorological databases (1986-93) acquired for the South Coast Air Basin .....	3-1
3-2	The fraction of days during the smog season with gaps in ozone data of two or more hours in length .....	3-3
3-3	The fraction of NO <sub>x</sub> data available for morning (4-10 am PST) observations for the summer months (May 1 - Oct. 31) for 1986-93 for the Coastal subregion .....	3-6
3-4	The fraction of NO <sub>x</sub> data available for morning (4-10 am PST) observations for summer months (May 1 - Oct. 31) for 1986-93 for the Metropolitan subregion .....	3-7
4-1	Subregions of the SoCAB employed in this study and air monitoring stations within each subregion for which air quality data were employed .....	4-3
4-2	Correlation coefficients between daily Basin ozone maximum concentration and average morning NO <sub>2</sub> and NO <sub>x</sub> concentrations in the Coastal and Metropolitan subregions .....	4-5

4-3	Correlation coefficients between daily Basin ozone maximum concentration and average morning NO <sub>2</sub> and NO <sub>x</sub> concentrations in the San Gabriel subregion	4-6
4-4	Correlation coefficients between daily Basin ozone maximum concentration and average morning NO <sub>2</sub> and NO <sub>x</sub> concentrations in the Inland Valley subregion	4-7
4-5	Correlation coefficients between daily Coastal/Metropolitan ozone maximum concentration and average morning NO <sub>2</sub> and NO <sub>x</sub> concentrations in the Coastal/Metropolitan subregion	4-10
4-6	Correlation coefficients between daily San Gabriel Valley ozone maximum concentration and average morning NO <sub>2</sub> and NO <sub>x</sub> concentrations in the San Gabriel Valley subregion	4-11
4-7	Correlation coefficients between daily Inland Valley ozone maximum concentration and average morning NO <sub>2</sub> and NO <sub>x</sub> concentrations in the Inland Valley subregion	4-13
4-8	Correlation coefficients between Tuesday evening (6-9 PDT) NO <sub>x</sub> concentration and Wednesday ozone maximum	4-14
4-9	Correlation coefficients between Friday evening (6-9 PDT) NO <sub>x</sub> concentration and Saturday ozone maximum	4-15
4-10	Correlation of previous evening's (6-9 PDT) San Gabriel Valley subregion NO <sub>2</sub> and daily ozone maximum in the Basin and in the San Gabriel Valley	4-16
5-1	Vandenberg AFB/San Diego mean 850 mb temperatures for 0500 PDT (1200 GMT) and 1700 PDT (0000 GMT) for the 28 high ozone concentration days and the 28 middle ozone concentration days during the 1990-93 smog seasons	5-32
5-2	Vandenberg AFB and San Diego 850 mb resultant wind speeds and directions for 0500 PDT (1200 GMT) and 1700 PDT (0000 GMT) for the 28 high ozone concentration days and the 28 middle ozone concentration days during the 1990-93 smog seasons	5-35
5-3	The number of occurrences of the pattern numbers for the	

28 middle and 28 high ozone days	5-39
5-4 Distribution of smog season days by node of the CART decision tree for each year in the period 1986-93	5-44
5-5 Distribution of smog season days by node of the CART decision tree for each of the two four-year periods 1986-89 and 1990-93	5-49
5-6 Distribution of the 28 high ozone days and the 28 middle ozone days by node of the CART decision tree	5-50
6-1 Daily (hourly-average) maximum ozone concentration (May 1 - Oct. 31) for Azusa in 1986 (pphm)	6-3
6-2 Weekday/weekend differences in daily ozone maxima between period I (1986-89) and period II (1990-93) in subregions of the South Coast Air Basin	6-29
6-3 Trends in hours equal or above the ozone NAAQS of 12 pphm	6-36
6-4 The application of Horie's CART scheme to weekdays and weekend days during 1986-89 and 1990-93	6-41
6-5 Means and standard deviations of the ten highest hourly average ozone concentrations for 32 stations for the period 1986-1990, and for the period 1990-93	6-43
6-6 Percentage changes in the means of the ten highest daily ozone maximum concentrations between 1986-89 and 1990-93 by subregion	6-47
6-7 Ozone trends for weekdays (WD) and for weekend days (WE) at stations in the SoCAB	6-48
6-8 Weekday/weekend day differences in ozone concentrations at stations grouped by subregion in the SoCAB for 1986-89 and 1990-93	6-50
6-9 The highest ozone concentrations in the SoCAB for each year in the period 1986-93	6-51
6-10 Mean morning (5-8 am PST) concentrations of NO <sub>2</sub> , NO <sub>x</sub>	

	(in pphm), and $\text{NO}_2/\text{NO}_x$ ratios on selected weekdays (Wednesdays and Thursdays), Saturdays, Sundays, and Mondays during May-October 1986-89	6-57
6-11	Mean morning (5-8 am PST) concentrations of $\text{NO}_2$ , $\text{NO}_x$ (in pphm), and $\text{NO}_2/\text{NO}_x$ ratios on selected weekdays (Wednesdays and Thursdays), Saturdays, Sundays, and Mondays during May-October 1990-93	6-57
6-12	t-Test for morning (5-8 am PST) concentrations of $\text{NO}_2$ and $\text{NO}_x$ , and for $\text{NO}_2/\text{NO}_x$ ratios for differences between weekdays and weekend days and Mondays for (May-October) 1986-89	6-60
6-13	t-Test for morning (5-8 am PST) concentrations of $\text{NO}_2$ , $\text{NO}_x$ , and $\text{NO}_2/\text{NO}_x$ ratios for differences between weekdays and weekend days and Mondays for (May-October) 1990-93	6-60
6-14	Average morning (5-8 am PST) $\text{NO}_2$ concentration (pphm) for the top ten ozone days in the indicated periods	6-61
6-15	Average morning (5-8 am PST) $\text{NO}_x$ concentration (pphm) for the top ten ozone days in the indicated periods	6-62
6-16	Mean 5-8 am (PST) $\text{NO}_2$ concentrations (pphm) for 28 high and 28 middle ozone days	6-64

## LIST OF FIGURES

<u>Figure Number</u>	<u>Title</u>	<u>Page</u>
1-1	Changes in means of the ten highest daily ozone concentrations (ppb) between 1986-89 and 1990-93 by station within the SoCAB . . .	1-19
1-2	Isopleth diagram for change in ambient 5-8 am (PST) NO <sub>x</sub> concentration (ppb) from 1986-89 to 1990-93 for top ten O <sub>3</sub> days in those periods . . . . .	1-20
4-1	Major portion of the SoCAB, with locations of air quality monitoring stations and subregions employed in this study, indicated . . . . .	4-2
4-2	Sunday Basin ozone maximum vs. Sunday morning Inland Valley NO <sub>2</sub> for Period I (1986-89) . . . . .	4-8
4-3	Friday San Gabriel Valley ozone maximum and Friday morning San Gabriel Valley NO <sub>2</sub> for period I (1986-89) . . . . .	4-12
5-1	0900-1000 PDT resultant winds for 28 high ozone days . . . . .	5-3
5-2	0900-1000 PDT resultant winds for 28 middle ozone days . . . . .	5-4
5-3	1200-1300 PDT resultant winds for 28 high ozone days . . . . .	5-5
5-4	1200-1300 PDT resultant winds for 28 middle ozone days . . . . .	5-6
5-5	1400-1500 PDT resultant winds for 28 high ozone days . . . . .	5-8
5-6	1400-1500 PDT resultant winds for 28 middle ozone days . . . . .	5-9
5-7	1600-1700 PDT resultant winds for 28 high ozone days . . . . .	5-10
5-8	1600-1700 PDT resultant winds for 28 middle ozone days . . . . .	5-11
5-9	Mean initial time of occurrence (PDT) of peak zone for 28 high high ozone days . . . . .	5-12

5-10	Mean initial time of occurrence (PDT) of peak ozone for 28 middle ozone days	5-14
5-11	Mean initial time of occurrence (PDT) of peak ozone for subset of high ozone days (N=19) on which peak ozone value within the SoCAB occurred in Glendora	5-16
5-12	1400-1500 PDT resultant winds for subset (N=19) of high ozone days on which Glendora reported the highest SoCAB ozone value	5-17
5-13	Mean initial time of occurrence (PDT) of peak ozone for subset of high ozone days (N=6) on which peak ozone value within the SoCAB occurred at Crestline	5-18
5-14	1400-1500 PDT resultant winds for subset (N=6) of high ozone days on which Crestline reported the highest SoCAB ozone value	5-20
5-15	Backward trajectories based on hourly resultant winds at SoCAB stations for Glendora high ozone days and Crestline high ozone days	5-22
5-16	West Los Angeles and Glendora wind roses for 0800-0900 PDT for high ozone concentration days and middle ozone concentration days during the 1990-93 smog seasons	5-24
5-17	West Los Angeles and Glendora wind roses for 1400-1500 PDT for high ozone concentration days and middle ozone concentration days during the 1990-93 smog seasons	5-26
5-18	Mean daily maximum surface temperature (°F) for 28 high ozone days	5-28
5-19	Mean daily maximum surface temperature (°F) for 28 middle ozone days	5-29
5-20	Map of California, indicating area of SoCAB	5-30
5-21	Distribution of mean (San Diego + Vandenberg AFB) 0500 PDT (1200 GMT) 850 mb temperatures for both 27 (out of 28) high ozone days (hatched) and 28 middle ozone days (solid)	5-33



5-22	Distribution of mean (San Diego + Vandenberg AFB) 1700 PDT (0000 GMT) 850 mb temperatures for both 26 (out of 28) high ozone days (hatched) and 28 middle ozone days (solid) . . . . .	5-34
5-23	Vandenburg and San Diego wind roses at 850 mb for 1700 PDT (0000 GMT) for 28 high ozone concentration days and 28 middle ozone concentration days during the 1990-93 smog seasons . . . . .	5-36
5-24	500 mb flow patterns defined by Cassmassi (1987) for ozone prediction stratification . . . . .	5-38
5-25	CART decision tree for daily SoCAB maximum ozone concentrations (including those measured at Glendora) . . . . .	5-40
6-1	Distribution of daily ozone maxima (May 1 - Oct. 31, 1986) for Central Los Angeles . . . . .	6-4
6-2	Distribution of daily ozone maxima (May 1 - Oct. 31, 1987) for Central Los Angeles . . . . .	6-5
6-3	Distribution of daily ozone maxima (May 1 - Oct. 31, 1988) for Central Los Angeles . . . . .	6-6
6-4	Distribution of daily ozone maxima (May 1 - Oct. 31, 1989) for Central Los Angeles . . . . .	6-7
6-5	Distribution of daily ozone maxima (May 1 - Oct. 31, 1990) for Central Los Angeles . . . . .	6-8
6-6	Distribution of daily ozone maxima (May 1 - Oct. 31, 1991) for Central Los Angeles . . . . .	6-9
6-7	Distribution of daily ozone maxima (May 1 - Oct. 31, 1992) for Central Los Angeles . . . . .	6-10
6-8	Distribution of daily ozone maxima (May 1 - Oct. 31, 1993) for Central Los Angeles . . . . .	6-11
6-9	Distribution of daily ozone maxima (May 1 - Oct. 31, 1986) for Azusa . . . . .	6-12

6-10	Distribution of daily ozone maxima (May 1 - Oct. 31, 1987) for Azusa	6-13
6-11	Distribution of daily ozone maxima (May 1 - Oct. 31, 1988) for Azusa	6-14
6-12	Distribution of daily ozone maxima (May 1 - Oct. 31, 1989) for Azusa	6-15
6-13	Distribution of daily ozone maxima (May 1 - Oct. 31, 1990) for Azusa	6-16
6-14	Distribution of daily ozone maxima (May 1 - Oct. 31, 1991) for Azusa	6-17
6-15	Distribution of daily ozone maxima (May 1 - Oct. 31, 1992) for Azusa	6-18
6-16	Distribution of daily ozone maxima (May 1 - Oct. 31, 1993) for Azusa	6-19
6-17	Distribution of daily ozone maxima (May 1 - Oct. 31, 1986) for Riverside	6-20
6-18	Distribution of daily ozone maxima (May 1 - Oct. 31, 1987) for Riverside	6-21
6-19	Distribution of daily ozone maxima (May 1 - Oct. 31, 1988) for Riverside	6-22
6-20	Distribution of daily ozone maxima (May 1 - Oct. 31, 1989) for Riverside	6-23
6-21	Distribution of daily ozone maxima (May 1 - Oct. 31, 1990) for Riverside	6-24
6-22	Distribution of daily ozone maxima (May 1 - Oct. 31, 1991) for Riverside	6-25
6-23	Distribution of daily ozone maxima (May 1 - Oct. 31, 1992) for Riverside	6-26
6-24	Distribution of daily ozone maxima (May 1 - Oct. 31, 1993)	

	for Riverside	6-27
6-25	Total number of ozone first stage alerts for the period 1986-93 (May 1 - Oct. 31) for Central Los Angeles, by day of the week	6-30
6-26	Total number of ozone first stage alerts for the period 1986-93 (May 1 - Oct. 31) for Azusa, by day of the week	6-31
6-27	Total number of ozone first stage alerts for the period 1986-93 (May 1 - Oct. 31) for Riverside	6-32
6-28	Number of ozone first stage alerts (May 1 - Oct. 31) for Central Los Angeles by day of the week for each individual year between 1986 and 1993	6-33
6-29	Number of ozone first stage alerts (May 1 - Oct. 31) for Azusa by day of the week for each individual year between 1986 and 1993	6-34
6-30	Number of ozone first stage alerts (May 1 - Oct. 31) for Riverside by day of the week for each individual year between 1986 and 1993	6-35
6-31	Cumulative ozone exposure for concentrations above 12 pphm (May 1 - Oct. 31) for Central Los Angeles, by day of the week for each year between 1986 and 1993	6-37
6-32	Cumulative ozone exposure for concentrations above 12 pphm (May 1 - Oct. 31) for Azusa, by day of the week for each year between 1986 and 1993	6-38
6-33	Cumulative ozone exposure above 12 pphm (May 1 - Oct. 31) for Riverside, by day of the week for each year between 1986 and 1993	6-39
6-34	Changes in means of the ten highest daily ozone concentrations (ppb) between 1986-89 and 1990-93 by station within the SoCAB	6-45
6-35	Isopleth diagram for change in ambient 5-8 am (PST) NO <sub>x</sub> concentration (ppb) from 1986-89 to 1990-93 for top ten O <sub>3</sub> days in those periods	6-46

7-1	0900-1000 PDT hourly-average surface winds for 28 August 1987	7-3
7-2	1200-1300 PDT hourly-average surface winds for 28 August 1987	7-5
7-3	1600-1700 PDT hourly-average surface winds for 28 August 1987	7-6
7-4	Initial times of occurrence (PDT) of peak ozone for 28 August 1987	7-7
7-5	Maximum hourly-average surface temperatures (°F) for 28 August 1987	7-9

## **GLOSSARY OF TERMS, ABBREVIATIONS, AND SYMBOLS**

AFB	Air Force Base
AQMD	Air Quality Management District
AQMP	Air Quality Management Plan
ARB	Air Resources Board
CART	Classification and Regression Tree
DI	Deviation Index
FAA	Federal Aeronautical Association
GEMPAK	General Meteorology Package
GMT	Greenwich Mean Time
IV	Inland Valley
LAX	Los Angeles International Airport
MDL	minimum detectable level
MM5	NCAR Mesoscale Model Version 5
mph	miles per hour
NCAR	National Center for Atmospheric Research
NMC	National Meteorological Center
NMHC	non-methanated hydrocarbon
NO <sub>2</sub>	nitrogen dioxide
NO <sub>x</sub>	nitrogen oxides
PM-10	particulate matter less than 10 µm in diameter

PDT	Pacific Daylight Time
PST	Pacific Standard Time
ROG	reactive organic gases
SCAQMD	South Coast Air Quality Management District
SCAQS	Southern California Air Quality Study
SGV	San Gabriel Valley
SoCAB	South Coast Air Basin
TPD	ton per day
TSP	total suspended particulate
UAM	Urban Airshed Model
UCLA	University of California, Los Angeles
VOC	volatile organic compound
WD	weekday
WE	weekend